

Helping Hands, Watchful Eyes; The Role of Home Visiting in Suicide Prevention

Maine Suicide Prevention Program In Partnership with NAMI Maine and Maine Medical Association

Education, Resources and Support—It's Up to All of Us.



Maine Suicide Prevention Program

A program of the Maine Center for Disease Control and Prevention since 1998

Statewide Activities Include:

- Crisis Hotline 1-888-568-1112
- Data collection, analysis & dissemination
- Dissemination of print materials through Substance Abuse and Mental Health Service's Information and Resource Center
800-498-0027 maineosa.org (8:00 a.m.—8:00 p.m., M – F)
- Training on suicide prevention and assessment
- Technical Assistance
- Annual Beyond the Basics conference



Suicide in the US, 2015


- 44,193 Americans died by suicide; about 1 person every 11.9 minutes
- Suicide deaths 2.5 times the number of homicides (homicides=17,793)
- 10th leading cause of death across the lifespan
 - 2nd leading cause of death for 15-34 year olds
- Men account for 77% suicides
 - 3 Female attempts per male attempt
- Veterans account for 20% of suicides
- Since 2009, suicides have exceeded motor vehicle crash related deaths

Suicide in Maine, 2013-2015

- 9th leading cause of death among all ages (previously 10th, 2012-2014)
- 2nd leading cause of death ages 10-34
- 4th leading cause of death ages 35-54
- Suicide deaths 9x homicide deaths
- Every 1.6 days someone dies by suicide
- Every 2 weeks a young person dies (ages 10-24)
- 233 suicide deaths per year on average
- Firearms most prevalent method of suicide (54%)




Age-Adjusted Suicide Death Rates: Maine, the Northeast and United States, 2000-2015




Data source: US CDC WISQARS Fatal Injury Data, National Vital Statistics System (NVSS)

Average Annual Suicide Deaths & Age-Adjusted Death Rates, by Age & Sex, Maine, 2012-2014



Data source: US CDC WISQARS Fatal Injury Data, National Vital Statistics System (NVSS); Rates for females ages 65-74, 75-84 suppressed due to small number of deaths. *Rate for men ages 85+ based on fewer than 10 deaths and should be interpreted with caution.

Lifespan Suicide Death Rates, by Age, US, 2014 (10 year age groups)



Suicide Among Older Adults

- Highest rate of any age group (for men)
- 87.5% of elder suicides in Maine are male (2013-15)
- 2013-15 Rates in Maine (17 per 100K)
 - -women 4.00 per 100,000
 - - men 34.01 per 100,000
- After age 60 rate declines for women
- Firearms most common means
- 66%-90% have diagnosable mental illness
- 2-4% completed suicides are terminally ill

Attempted Suicides- A Call For Help!



Characteristics of Elderly Suicide Attempts

Ask about a history of attempts!

- **More secretive:** Fewer warnings of intent
- **More planful:** Attempts are more planned, determined
2/3 have high suicide intent scores
- **More lethal**
 - Less likely to survive a suicide attempt due to use of more violent and immediate methods
 - Also more frail

Conwell Y, Duberstein PR, Cox C, Hermann J, Forbes N, & Caine ED. Age differences in behaviors leading to completed suicide. *American Journal of Geriatric Psychiatry*. 1998;6(2):122-6.

10

Discussion

Overall, suicide rates among older adults have fallen since 1930.
What changes in policy, supports, cultural attitudes and healthcare practices have supported this trend?

What do you see as priorities that would support reduction of suicide rates among older men?

Suicide in Older Adults

Clarification of Attitudes



Examining Our Own Attitudes

- What associations do we have to the word “suicide”?
- What do we “know” about suicide?
- How has suicide impacted your life?
- What do we “know” about people who are suicidal?

Values Clarification

- Is there a difference between an adolescent suicide and an older adult suicide?
- For someone diagnosed with a terminal illness is it still a suicide?
- What is the difference between “death with dignity” and suicide?
- Is there such a thing as “rational suicide”?

What is the message?



Warning Signs Risk Factors Protective Factors

Risk Factors among Older Adults

- **Male**, white and old old (esp. after losses)
- **Depression** (esp. untreated),
- Prior suicide attempts,
- Marked feelings of **hopelessness**,
- Co-morbid medical conditions limiting functioning,
- Pain and **declining role function**,
- Social/familial **isolation/cut-offs or losses**
- Rigid inflexible personality
- Access to lethal means (esp. **firearms**)
- Substance abuse

Men as a High Risk Group

- 80% of suicides
- Gender disparity highest in elders (especially white)
- Gender issues include:
 - Poor help-seeking
 - Men less likely to talk to someone
 - Difficulty recognizing and expressing emotions
 - Increased substance abuse
 - Use more lethal means
 - Feeling like a burden
 - Struggle between belongingness and independence

Warning Signs

What have you seen that tells you that a person is at increased risk?

- In your center?

- In the community/home?



Clear Signs Of A Suicidal Crisis

1. Someone threatening to hurt or kill themselves
2. Someone looking for the means (gun, pills, rope etc.) to kill themselves
3. Someone showing clear distress/ agitation/ anxiety

Get the facts and take action!

Call **911** if lethal means is present
Call **Crisis Hotline** if no means present

Warning Signs of Suicide in Elders

- Direct or indirect communication
 - Hopelessness, Purposelessness, Isolation,
- Giving away possessions
- Getting affairs in order
- Saying good bye
- Sudden interest or disinterest in religion (change in interest)
- A specific plan for how they will die

Warning Signs: Depression

Physical

- Aches, pains, or physical complaints
- Marked changes in appetite
- Change in sleep patterns
- Fatigue

Emotional

- Pervasive sadness
- Apathy
- Decreased pleasure
- Crying for no apparent reason
- Indifference to others

Changes in Thoughts and Feelings

- Feelings of hopelessness and helplessness
- Feelings of worthlessness
- Impaired concentration
- Problems with memory
- Indecisiveness
- Recurrent thoughts of death and suicide

Changes in Behavior

- Loss of interest in previously enjoyed activities
- Neglect of personal appearance
- Withdrawal from people
- Increased use of alcohol
- Increased agitation / anxiety
- Talking about the "end"

Adapted from Schmall V, Lawson L, Stehl R. Depression in Later Life: Recognition and Treatment. Pacific Northwest Extension publication. Corvallis, Ore, 1993

Warning Signs



Protective Factors

- ☑ **Skills** to think, communicate, solve problems, manage anger and other negative emotions,
- ☑ **Purpose & value** in life-hope for future, pets, life focus...
- ☑ **Personal characteristics**- health, positive outlook, spirituality or religious belief
- ☑ **Supports**- friends, family, and other caring people, health care access, transportation
- ☑ **Safe Environment** – restricted access to lethal means

From a Suicidal Person's Point of View

- **Crisis point** has been reached
- **Pain** is unbearable
- **Solutions** to problems seem unavailable
- **Thinking** is affected
- **HOWEVER:**
- **Ambivalence** exists
- **Communicating** distress is common
- **Invitations** to help are often extended
 - Less often or open for older adults

How to Help?

Intervention

- It all starts with a conversation
- Active intervention is needed
- Engagement is essential
- Importance of connections/ **breaking isolation**
- Reduce the level of risk by removing all **lethal means**
- **Invitations** are often extended to people based on fit
- **Invitations** are often extended to people based on opportunity and availability

What IS Helpful

- 1) **Show You Care**—Listen carefully—Be genuine

"I'm concerned about you . . . about how you feel."

- 2) **Ask the Question**—Be direct, caring and non-confrontational

"Are you thinking about ending your life?"

- 3) **Get Help**—Do not leave him/her alone

"You're not alone. Let me help you."

Resources for Help

What are YOUR resources?



Resources for Help

To address the Crisis

- 911 or Law enforcement
- **Statewide Crisis Hotline (888-568-1112)**
- Local Crisis Agency, Mental Health Clinicians and Facilities
- Hospital emergency room staff or PCP office/rural health center in rural areas

For follow-up, support & information after the crisis

- Private counselors/therapist
- Faith Community
- Local Health Center
- 211
- Maine's Intentional Warmline: 1-866-771-9276

When to Call Crisis

- Crisis clinicians are:
 - Available 24 / 7
 - Clinicians can often come to your location for an assessment
- Call for a phone consult when you are:
 - Concerned about someone's mental health
 - Need advice about how to help someone in distress
 - Worried about someone and need another opinion
- The phone call is free

1-888-568-1112

Crisis Intervention Teams

The Crisis Intervention Team program trains police, correctional officers and first responders about mental illness and methods to deal with mental health emergency and crisis situations safely.

But it is not just a training, CIT transforms how the entire community responds to psychiatric crisis by creating an ongoing collaboration that supports jail diversion



If you need to call the police for a mental health emergency, ask for a CIT trained officer

Key Actions For Healthcare Providers

- Routine standard screening for depression,
- Use collaborative Tx of depression,
- Optimize treatment of pain, anxiety... to address quality of life issues,
- Include collateral folks in treatment discussions
- Active management after a suicide attempt or crisis.
 - Means restriction and safety planning
 - Increased outreach, care management and follow-up
 - Referrals for community programs

Key Actions for Aging Service Providers

- Training for staff on Warning signs and Risk factors and intervention skills
- Depression screening in non-clinical and community settings
- Center-based social programs
- **Outreach, outreach, outreach**
 - Target isolation
 - Activate family and social; supports
 - Meals on Wheels
 - Home visiting
 - Mail carriers, Faith community, Home handyman services...
 - Other?...

Survivors of Suicide



Paul A. LePage, Governor Mary C. Mayhew, Commissioner

Effect of Suicide

- **The Loss is:**
 - Sudden
 - Unexpected
 - Premature
 - Self-inflicted
- **The Reaction is:**
 - Shock, hurt, anger
 - Loss and grief
 - Questions & torment
 - Guilt and regret



Survivors of Suicide

- Struggle to make meaning of the loss
- Suffer from overwhelmingly complicated feelings
- May take a long time to grieve
- Need understanding and support
- Youth survivors have special issues

How YOU can be supportive after a suicide

- Acknowledge the loss
- Use the name of the deceased
- Share your presence
- Share a special memory/story
- Acknowledge the good things
- Stay in touch
- Recommend grief counseling or grief support groups

Take Care of Yourself

- Acknowledge the intensity of your feelings
- Seek support from others, de-brief
- Share your feelings with family/friends
- Avoid over – involvement. Never act in isolation
- Know that you are not responsible for another person's choice to end their life

MSPP Training and Technical Assistance

- *Suicide Prevention Gatekeeper Training*
- *Suicide Prevention: Training of Trainers*
 - *Supports capacity to offer Awareness Sessions*
- *Suicide Prevention Protocol Development Training & TA*
- *Suicide Assessment for Clinicians*

Contact NAMI Maine Suicide Prevention Training Coordinator for more details
mspp@namimaine.org

Contact: For Additional Support

Greg A Marley, LCSW 207-622-5767 ext. 2302
gmarley@namimaine.org

Training Program Inquiries 207-622-5767 MSPP@namimaine.org

Dee Kerry 620-0806 Maine Medical Association
dkerry@mainemed.com

Sheila Nelson, MSPP Program Coordinator 207-287-3856
Sheila.Nelson@maine.gov

Resources

- See Handout

Before you leave. . .

Any Questions?



**Thank you for learning about
suicide prevention. . .**



Before you leave. . .

The Department of Health and Human Services (DHHS) does not discriminate on the basis of disability, race, color, creed, gender, sexual orientation, age, or national origin, in admission to, access to, or operations of its programs, services, or activities, or its hiring or employment practices. This notice is provided as required by Title II of the Americans with Disabilities Act of 1990 and in accordance with the Civil Rights Act of 1964 as amended, Section 504 of the Rehabilitation Act of 1973, as amended, the Age Discrimination Act of 1975, Title IX of the Education Amendments of 1972 and the Maine Human Rights Act and Executive Order Regarding State of Maine Contracts for Services. Questions, concerns, complaints or requests for additional information regarding the ADA may be forwarded to DHHS' ADA Compliance/EEO Coordinators, 11 State House Station - 221 State Street, Augusta, Maine 04333, 207-287-4289 (V), 207-287-3488 (VI). TTY users call Maine relay 711. Individuals who need auxiliary aids for effective communication in program and services of DHHS are invited to make their needs and preferences known to the ADA Compliance/EEO Coordinators. This notice is available in alternate formats, upon request.

