

**MSAT Program Emergency Cardiac Care (CPR), First Aid,  
and Immunization Requirements**  
Updated August 2026

**Proof of current certification in Emergency Cardiac Care\* that includes a minimum of the following:**

- a. Adult & Pediatric CPR
- b. AED
- c. 2<sup>nd</sup> Rescuer CPR
- d. Airway Obstruction
- e. Barrier Devices (e.g., pocket mask, bag valve mask)

\*Courses that meet the minimum requirement include, but are not limited to American Heart Association ACLS, American Heart Association BLS Healthcare Provider, and American Red Cross CPR/AED for the Professional Rescuer. Further information regarding CPR requirements can be found on the [Board of Certification Emergency Cardiac Care website](#).

**Proof of current certification in Basic First Aid\* that includes a minimum of the following:**

- a. Burns & Wounds
- b. Splinting & Bone/Joint Injuries
- c. Sudden Illness
- d. Shock
- e. Controlling Bleeding
- f. Head/Neck/Back Injuries
- g. Heat/Cold Emergencies

\*Courses that meet the minimum requirement include but are not limited to American Red Cross First Aid Basics.

**Proof of current immunizations as follows:**

1. Three (3) dose series **OR** (2) HepB vaccinations required **AND** Positive Surface Antibody titer lab report **OR** any document indicating titer results is required
  - Antigen test will not be accepted
  - If titer is negative/low/equivocal, must repeat 3 dose or 2 dose HepB series
  - Repeat TITER is not required
2. Two (2) doses of MMR vaccination **OR** a positive titer (lab report required).
  - If titer is negative/low/equivocal, then 2 administrations of the vaccine or a booster dose as recommended by healthcare provider must be provided, and no further testing will be required.
3. Two (2) doses of Meningococcal vaccination are required
4. Tdap within 10 years **OR** A previous Tdap dose and a current Td booster within 10 years.
  - Td booster is NOT ACCEPTED
5. Two (2) doses of Varicella vaccination **OR** a positive titer (lab report required)
  - If titer is negative/low/equivocal, then 2 administrations of the vaccine or a booster dose as recommended by healthcare provider must be provided, and no further testing will be required